

LETTER TO THE EDITOR

A class of parameter-dependent commuting matrices**B Sriram Shastry**

Department of Physics, University of California, Santa Cruz, CA 95064, USA

Received 21 March 2005, in final form 29 April 2005

Published 25 May 2005

Online at stacks.iop.org/JPhysA/38/L431**Abstract**

We present a novel class of real symmetric matrices in arbitrary dimension d , linearly dependent on a parameter x . The matrix elements satisfy a set of nontrivial constraints that arise from asking for commutation of pairs of such matrices for all x , and an intuitive sufficiency condition for the solvability of certain linear equations that arise therefrom. This class of matrices generically violates the Wigner von Neumann non-crossing rule, and is argued to be intimately connected with finite-dimensional Hamiltonians of quantum integrable systems.

PACS numbers: 71.10.Fd, 02.30.Ik, 02.10.Yn

Introduction

The study of linearly parameter dependent matrices is very popular in the context of quantum chaos [1–4], where it models the change in universality class of quantum systems, or the temporal variation of correlations within the same class via the matrix Brownian motion model of Dyson [5–7]. The present work has a different goal, that of formulating constraints so that the system remains ‘regular’ despite mixing with another matrix. In fact, such linearly parameter dependent matrices are prototypes of quantum integrable models. The well-known case of the anisotropic Heisenberg model solved by Bethe’s ansatz [8] possesses ‘infinite’ higher conservation laws, *every one of them being linear in the anisotropy parameter* [9]¹.

The immediate motivation for our work comes from a study of the parameter dependence of eigenvalues of blocks of finite-dimensional matrices arising from completely integrable models of interacting quantum systems, such as the Heisenberg model [8] or the 1D Hubbard model [10]. The famous Wigner–von Neumann (WvN) [11] non-crossing rule for parametric evolution of eigenvalues in quantum mechanics is a fundamental guiding principle for understanding level crossings, with violations or exceptions being termed as ‘accidental degeneracies’. The term ‘accident’ is used since usually there is no specific ‘spacetime symmetry’ responsible for such a degeneracy. There is a general belief that *dynamical*

¹ The Hubbard model’s higher conservation laws on the other hand [15] involve a polynomial of degree ≥ 1 in x (or U).

symmetries, i.e. operators *dependent on the interaction parameter* also lead to degeneracies and violations of the WvN rules, despite the lack of a general group theoretic argument of the type that spacetime symmetries allow².

Dynamical symmetries occur in most integrable systems. Our interest is also in sharpening the notion of complete or exact integrability in the context of finite-dimensional systems. In classical mechanics we have a very clear statement about the meaning of complete integrability, namely that the number of degrees of freedom equals the number of conservation laws that are mutually consistent. The meaning of the term ‘degrees of freedom’ is quite unambiguous. For example in the trivial case of a set of harmonically coupled oscillators, it is the number of oscillators so coupled. In quantum mechanics, we do have similar unambiguous models, such as the Calogero–Sutherland–Moser system [12] in the continuum, or the Toda [13] lattice, where there exists a natural definition of a degree of freedom, in complete parallel to the classical situation, essentially the number of ‘particle’-type variables. However, in the case of other quantum integrable models that arise in condensed matter physics, such as the Heisenberg spin chain or the 1D Hubbard model, the situation is ambiguous: the number of spin flips, or particles is variable. These models are defined on a discrete lattice, and have a state space that is *in general finite*, and lead to finite-dimensional matrices depending upon a parameter, say the spin space anisotropy or the interaction strength. Presented with a finite-dimensional matrix arising by restricting such a model to a finite lattice, it is challenging to distinguish it from other matrices of the same dimension. Without referring back to the defining parent models, it is generally impossible to recognize their being ‘integrable’, whatever that word implies! Indeed, an extreme and skeptical view would challenge the very notion of complete integrability in a finite-dimensional setting. One may argue that there are always d commuting independent matrices for *any given matrix* H ; one merely diagonalizes the matrix and in its eigenbasis, constructs the d projection operators $P_j = |j\rangle\langle j|$, so that $[H, P_j] = 0 = [P_i, P_j]$.

Our viewpoint in this work is that the *parameter dependence of eigenvalues contains the essence of quantum integrability* for such finite-dimensional models. These lead to a violation of the WvN rules, and hence to Poisson statistics of the energy level separation. While the relationship between dynamical symmetries and level crossings is not yet precisely established, there are several studies that indicate a deep relationship between them. For example, the beautiful numerical work of Heilmann and Lieb in 1971 [14] on the one-dimensional Hubbard model on a six-site ring, shows that after all the known spacetime and internal space symmetries are carefully extracted, the finite-dimensional blocks of matrices labelled by the appropriate quantum numbers, displaying scant regard for the WvN rules, have a large number of level crossings as the interaction constant is varied. More recent work of Yuzbashyan *et al* [15] has examined the detailed connection between these level crossings and the dynamical symmetry operators of the Hubbard model found in 1986 by Shastry [16], and provided considerable insight into this phenomenon. In particular, there is an explicit algebraic demonstration for $d = 3$ that dynamical symmetries definitely imply level crossings.

Another related and prominent manifestation of integrability is that the energy level statistics of such models display Poisson-type behaviour that allows spacings to be arbitrarily small [17]. This is in sharp contrast with the level spacings of generic (i.e. nonintegrable) models that exhibit level repulsion as expected from the WvN rule, and follow one of the

² The Runge Lenz vector for the hydrogen atom is a celebrated example of a dynamical symmetry in quantum theory, giving an explanation of the ‘permanent degeneracy’ between different $|\bar{L}|$ eigenstates. It is also untypical in that it can be cast into a group theoretic framework, leading to $SO(4)$ dynamical symmetry group of that problem. Our concern below is more with level crossings and their relationship with dynamical symmetries, where such a group structure is not available.

three typical behaviours relevant to their class of quantum systems—namely the Gaussian orthogonal, unitary or symplectic classes [18].

It remains, however, to state explicit conditions on matrices in arbitrary finite dimensions, which could identify the proclivity for level crossings, and hence presumably define completely integrable cases without reference to a parent quantum model. This goal is achieved in this work, we present a set of conditions, and a class of matrices satisfying them in any dimension d . We find through examples that this class of matrices automatically leads to Poisson-type statistics for the energy level separation, and also an abundance of level crossings.

Our main results follow from asking for the conditions for two parameter-dependent matrices $\alpha(x)$ and $\beta(x)$ to commute with each other for all values of the parameter. Upon imposing a very intuitive sufficiency condition of autonomy (type I matrices as explained below), we find linear as well as nonlinear constraints, so that $\alpha(x)$ has $3d - 1$ independently assignable real parameters. For a given $\alpha(x)$ we have the freedom of $d + 1$ parameters in the $\beta(x)$ matrices, all of which are of type I and display level crossings as x is varied.

The matrix equations

We consider real symmetric matrices in d dimensions, linearly dependent on a real parameter x through

$$\alpha(x) = a + xA, \quad (1)$$

where a is a diagonal matrix with diagonal entries $\{a_1, a_2, \dots, a_d\}$ and A is a real symmetric matrix³. In the matrix α we have $d(d - 1)/2$ off-diagonal variables $A_{i,j}$, d variables a_j and a further d variables $A_j \equiv A_{j,j}$, in addition to the real variable x . In an identical fashion we consider another matrix

$$\beta(x) = b + xB, \quad (2)$$

where the diagonal matrix b has entries $\{b_1, b_2, \dots, b_d\}$ and B is another real symmetric matrix. Clearly at $x = 0$ the matrices α and β commute. We now ask the question, what are the conditions under which these commute for *arbitrary values of x* ? This clearly leads to two independent conditions

$$[a, B] = [b, A] \quad \text{and} \quad [A, B] = 0. \quad (3)$$

The first set of $d(d - 1)/2$ conditions is expressible in terms of an antisymmetric matrix $S_{i,j}$ as follows:

$$S_{i,j} = \frac{A_{i,j}}{a_i - a_j} = \frac{B_{i,j}}{b_i - b_j} \quad \text{for } i < j. \quad (4)$$

The second set of equations can be written compactly using equation (4) in terms of

$$Y_{i,j}[\alpha] \equiv p_{i,j} - \frac{1}{S_{i,j}(a_i - a_j)} \sum_{l \neq i,j} S_{i,l} S_{l,j} (a_l - a_j) \quad (5)$$

where

$$p_{i,j} = \frac{A_i - A_j}{a_i - a_j}.$$

The $d(d - 1)/2$ equations (3) can be written as

$$Y_{i,j}[\alpha] = Y_{i,j}[\beta], \quad (6)$$

³ It is clear that the commutators of interest are unchanged by adding a constant times the identity matrix to α or β , the equations that follow reflect this invariance.

where $Y_{i,j}[\beta]$ is obtained from the same formula (5), with $(a_j, A_j) \leftrightarrow (b_j, B_j)$. Although it is possible to symmetrize $Y_{i,j}$ in i and j , for future use it is better to use the present unsymmetric form. As expected, equation (6) is symmetric in α and β . At this point, we disturb this symmetry, and rearrange terms so that we can solve for β if α were given. This implies with $(b \wedge a)_{i,j,l} \equiv [(b_i a_j - b_j a_i) + (b_j a_l - b_l a_j) + (b_l a_i - b_i a_l)]$

$$\xi_{i,j} \equiv B_i - B_j = p_{i,j}(b_i - b_j) + \frac{1}{a_i - a_j} \sum_{l \neq i,j} \frac{S_{i,l} S_{l,j}}{S_{i,j}} (b \wedge a)_{i,j,l}. \quad (7)$$

This set of equations is linear in the $2d$ variables $\{b_1, b_2, \dots, b_d\}$ and $\{B_1, B_2, B_3, \dots, B_d\}$, and provides conditions for the matrix $\beta(x)$ to commute with $\alpha(x)$ for all x . The off-diagonals of the matrix B namely $B_{i,j}$ are already fixed by equation (4) once the b'_j s are picked in terms of the $S_{i,j}$. For a given $\alpha(x)$ then, we have the freedom of choosing these $2d$ variables subject to the $d(d-1)/2$ constraints (7), which for large d , are many more than the number of variables available. These constraints are best expressed in terms of a ‘triangle law’ for any three distinct indices i, j, k :

$$C(i, j, k) \equiv \xi_{i,j} + \xi_{j,k} + \xi_{k,i} = 0. \quad (8)$$

Note that C is antisymmetric under exchange of any pair of arguments. Since these constraints are linear in b'_j s we collect the coefficients and write

$$C(i, j, k) = \mu(i; j, k)b_i + \mu(j; k, i)b_j + \mu(k; i, j)b_k + \sum_{l \neq i,j,k} \nu(l; i, j, k)b_l. \quad (9)$$

μ is defined below in equation (19) and

$$\nu(l; i, j, k) = \frac{S_{i,l} S_{l,j}}{S_{i,j}} + \frac{S_{j,l} S_{l,k}}{S_{j,k}} + \frac{S_{k,l} S_{l,i}}{S_{k,i}}. \quad (10)$$

The solutions of equations (8) may be classified as being of two types. Type I solutions correspond to requiring the coefficients of every b_r in equation (9) to vanish individually. Such a choice is sufficient without being necessary, and gives us constraints on the S matrix all by itself. It also leads to *autonomous* constraints on α , i.e. constraints involving the variables $a_j, A_j, S_{i,j}$, but not the β variables. Type II solutions include all other possibilities, where the coefficients of b_r are not all vanishing. These include the trivial solution $\alpha = \beta$, and are less interesting as such. Type I solutions give us families of matrices α as we show below, which are definable in terms of relations involving S 's themselves, and between these and the remaining variables.

Continuing our study of type I solutions, we therefore equate the coefficients of b 's individually to zero, giving for all ${}^d C_3$ distinct triples i, j, k the three index formulae,

$$\mu(i; j, k) = 0 \text{ plus cyclic permutations of } i, j, k. \quad (11)$$

Likewise we get ${}^d C_4$ four index formulae for each distinct quadruple of indices, $\nu(l; i, j, k) = 0$, or rearranging a bit:

$$S_{i,l} S_{l,j} S_{j,k} S_{k,i} + S_{j,l} S_{l,k} S_{k,i} S_{i,j} + S_{k,l} S_{l,i} S_{i,j} S_{j,k} = 0. \quad (12)$$

Note that the expression on the LHS is *fully symmetric* in the four variables. Although this relation involves quartics in the S 's, inspection shows that for the case of real symmetric matrices, a considerable simplification occurs and this constraint can be written in terms of bilinears if one inverts the matrix elements (not the matrix itself!) of S and defines

$$R_{i,j} = \frac{1}{S_{i,j}}. \quad (13)$$

The vanishing of ν in equation (12) can be restated as the (fermionic Wick’s theorem type) requirement that the totally antisymmetric symbol ϕ vanishes for all distinct quadruples i, j, k, l

$$\phi_{(i,j,k,l)} \equiv R_{i,j}R_{k,l} - R_{i,k}R_{j,l} + R_{i,l}R_{j,k} = 0. \tag{14}$$

For large d , the four index constraints (14) are $\sim d^4/4!$ in number, representing a huge overdetermination since the number of variables available, namely the $S_{i,j}$ are only $\sim d^2/2$ in number⁴. Fortunately these identities are not all independent, and there exists an important extra identity relating a set of five indices that can be proved. For any distinct set of five indices i, j, k, l, m , we can easily see that

$$\phi_{(i,j,k,l)}R_{l,m} = -\phi_{(i,j,l,m)}R_{l,k} + \phi_{(i,k,l,m)}R_{l,j} - \phi_{(j,k,l,m)}R_{l,i}. \tag{15}$$

Using this five index identity we can show that the number of independent quadruple relations is only $(d - 2)(d - 3)/2$ in number. These may be chosen to be

$$\phi_{(1,2,3,4)} = 0, \phi_{(1,2,3,5)} = 0, \dots, \phi_{(1,2,d-1,d)} = 0. \tag{16}$$

From these equations, we can satisfy all others using equation (15) repeatedly. As an example consider $\phi_{(3,4,5,6)}$, we merely multiply by $R_{3,1}$ so that on using equation (15), we find

$$\phi_{(3,4,5,6)}R_{3,1} = \phi_{(3,1,5,6)}R_{3,4} - \phi_{(3,1,4,6)}R_{3,5} + \phi_{(3,1,4,5)}R_{3,6}, \tag{17}$$

and each of these can be similarly processed further, e.g.

$$\phi_{(1,3,4,6)}R_{1,2} = \phi_{(1,2,4,6)}R_{1,3} - \phi_{(1,2,3,6)}R_{1,4} + \phi_{(1,2,3,4)}R_{1,6} \tag{18}$$

and reduced to forms involving $\phi_{(1,2,l,m)}$ which vanish according to our list of equation (16).

We now turn to the study of $\mu(i; j, k)$ which is written compactly as

$$\mu(i; j, k) = Y_{i,j} - Y_{i,k} - \frac{S_{i,j}S_{i,k}}{S_{j,k}} \tag{19}$$

involving the same variable $Y_{i,j}$ that we encountered in equation (5).⁵ The function μ is antisymmetric in j, k as it stands. It can be antisymmetrized in all three variables by defining $\tilde{\mu}(i, j, k) \equiv (a_i - a_j)(a_i - a_k)\mu(i; j, k)$, which satisfies

$$\begin{aligned} \tilde{\mu}(i, j, k) = & (a_i - a_j) \left[A_k - \frac{A_{i,k}A_{j,k}}{A_{i,j}} \right] + (a_j - a_k) \left[A_i - \frac{A_{i,j}A_{i,k}}{A_{k,j}} \right] \\ & + (a_k - a_i) \left[A_j - \frac{A_{j,k}A_{k,i}}{A_{k,i}} \right] - (a_i - a_j)(a_i - a_k) \sum_{r \neq i,j,k} S_{i,r} \left(\frac{A_{j,r}}{A_{i,j}} - \frac{A_{k,r}}{A_{i,k}} \right). \end{aligned} \tag{20}$$

We comment on some important properties of these formulae. The first part of equation (20) consisting of three terms is explicitly antisymmetric in the three indices. The last line is antisymmetric in j, k but not manifestly so in i, j . We add to it a term with i, j exchanged, leading to

$$\tilde{\mu}(i, j, k) + \tilde{\mu}(j, i, k) = \sum_{r \neq i,j,k} (a_i - a_j)(a_k - a_r) \left[\frac{S_{i,r}S_{r,j}}{S_{i,j}} + \frac{S_{j,r}S_{r,k}}{S_{j,k}} + \frac{S_{i,r}S_{r,k}}{S_{k,i}} \right] \tag{21}$$

but it vanishes on using the four index formula (12), whereby $\tilde{\mu}(i, j, k)$ is antisymmetric in all three indices.

⁴ We observe that the argument presented here has a tactical parallel to that in the solution of most quantum integrable models, e.g. using Bethe’s ansatz; one has a greatly overdetermined system, and the crux of integrability lies in the Yang–Baxter-type consistency of such equations.

⁵ The α dependence of the $Y_{i,j}$ is suppressed for brevity.

One more beautiful property of μ is:

$$\mu(i; j, k) - \mu(i; j, l) = Y_{i,l} - Y_{i,k} + \frac{S_{i,j}S_{i,k}}{S_{k,j}} + \frac{S_{i,j}S_{i,l}}{S_{j,l}} = \mu(i; l, k), \quad (22)$$

where the last line follows again from the use of the four index formula (12). Thus the total number of independent constraints of the μ -type is only $(d - 2)$ in number (see footnote 4), instead of the apparently huge number dC_3 . We may choose them most simply as

$$\mu(1; 2, 3) = 0, \dots, \mu(1; 2, d - 1) = 0, \quad \mu(1; 2, d) = 0. \quad (23)$$

The reader can verify that all other constraints of the type $\mu(i, j, k) = 0$ are obtainable from these $d - 2$ by using equation (22) and the antisymmetry of $\mu(i, j, k)$.

Constraints, variables and consistency

We now recount the number of available variables versus the constraints and show how generic matrices of type I can be constructed. Firstly we construct the S matrix satisfying the $(d - 2)(d - 3)/2$ constraints (16): this can be done by assigning arbitrary values to $(2d - 3)$ matrix elements of S and computing the rest from the constraints. A particularly convenient choice is to assign values to $S_{1,j}$ and $S_{2,j}$ for $\{2, 3\} \leq j \leq d$, whereby the constraints (16) reduce to *linear equations* for the other matrix elements.

Having determined the S matrix, we next construct the α matrices. A straightforward strategy is to assign arbitrary values to the $d + 2$ variables $\{a_j\}$, A_1 , A_2 , and then using the $d - 2$ constraints (23) as linear equations for the remaining A_j . Thus the total number of freely assignable variables for constructing the α matrix is $3d - 1$.

We can next determine a β matrix by assigning arbitrary values to the set of $d + 1$ variables $\{b_j\}$ and one of the B_j 's (say B_1), and using the linear equations (7) for $\xi_{i,1}$ to determine the rest. Thus $d + 1$ is the number of independent β -type matrices of type I for a given α .

By construction the resulting β and α matrices commute for all x . It is further clear that equations (6) are satisfied identically. Hence it follows that $\mu(i; j, k) = 0$ for all triples is satisfied whether we use $Y_{i,j}[\alpha]$ or $Y_{i,j}[\beta]$. This guarantees that starting with a matrix $\alpha(x)$ of type I, the resulting matrix $\beta(x)$ is *automatically* of type I.

We noted that in addition to the solutions presented above, there are some beautiful special solutions of these constraints (14) belonging to the class of Toeplitz matrices. From inspection and using various addition theorems, it is clear that the following class of R 's satisfies this constraint identically:

$$R_{i,j} = (\kappa(i) - \kappa(j)) \quad \text{or} \quad R_{i,j} = \sin(\kappa(i) - \kappa(j)), \quad (24)$$

where $\kappa(j)$ is an arbitrary function of its argument.

We plan to return to the problem of Hermitean as well as symplectic matrices in a future work⁶. One expects similar results to the ones presented here, but with more elaborate constraints. The results in $d = 3$ [15] are contained in our present ones, and correspond to the simple case of requiring $\mu(1; 2, 3) = 0$; this result is sufficient to make the discriminant vanish at points in x . For higher d the analysis of the discriminant is more difficult; however, examination of several examples of type I matrices leads us to conjecture that these *always lead to level crossings* [19]. A direct algebraic proof involves examining the condition for vanishing of the discriminant of the matrices, and is currently being pursued.

⁶ In the case of Hermitean matrices, the analysis follows similar lines, and indeed until equation (12) the same equations are true, but with an anti-Hermitian S . However, it cannot be reduced to the form of equation (14), there are non-trivial phase factors that arise, and hence the constraints seem to be intrinsically *quartic*.

An important issue concerns *translation invariance* of the results for type I matrices. It is not obvious that a matrix of type I remains so if we shift x by a constant and absorb the change into the diagonal piece a by re-diagonalizing using a suitable orthogonal transformation. We have verified that the results *do possess* this translation invariance in the parameter x , both numerically (for small d) and analytically. A detailed proof using the parameter derivatives of the constraints (23), (12), using the Pechukas flow equations [20] is essentially complete, and will be published separately. In future work we hope to address several physical models using these constraints (and some obvious variants), to check for their compliance.

Acknowledgment

I acknowledge support from the NSF grant DMR 0408247.

References

- [1] Berry M V 1987 *Proc. R. Soc. A* **413** 183
- [2] Berry M V 1983 *Les Houches Lecture Notes XXXVI* ed R Hellman and G Iooss (Amsterdam: North Holland)
- [3] Bohigas O and Gianonni M J 1984 *Chaotic Motion and Random Matrix Theory (Springer Lecture Notes in Physics vol 209)* (Berlin: Springer) p 1
- [4] Guhr T and Muller-Groeling A 1977 *J. Math. Phys.* **38** 1870
- [5] Dyson F J 1962 *J. Math. Phys.* **3** 1191
- [6] Simon B D, Lee P A and Altshuler B I 1993 *Phys. Rev. Lett.* **70** 4122
- [7] Narayan O and Shastry B S 1993 *Phys. Rev. Lett.* **71** 2106
- [8] Bethe H A 1931 *Z. Phys.* **71** 205
- [9] Thacker H 1986 *Physica D* **18** 348
- [10] See, e.g., Montorosi A (ed) 1992 *The Hubbard Model* (Singapore: World Scientific)
- [11] Neumann J v and Wigner E 1929 *Phys. Z.* **30** 467
- [12] Calogero F 1969 *J. Math. Phys.* **10** 2191
Sutherland B 1971 *J. Math. Phys.* **12** 246
Moser J 1975 *Adv. Math.* **16** 197
- [13] Toda M 1967 *J. Phys. Soc. Japan* **23** 501
Sutherland B 1978 *Rocky Mountain J. Math.* **8** 413
Gutzwiller M 1981 *Ann. Phys.* **133** 304
- [14] Heilmann O J and Lieb E H 1971 *Ann. N. Acad. Sci.* **172** 583
- [15] Yuzbashyan E A, Altshuler B L and Shastry B S 2002 *J. Phys. A: Math. Gen.* **35** 7525
- [16] Shastry B S 1986 *Phys. Rev. Lett.* **56** 1529, 2453
Shastry B S 1988 *J. Stat. Phys.* **50** 57
- [17] Poilblanc D, Ziman T, Bellissard J, Mila F and Montambaux G 1993 *Europhys. Lett.* **22** 537 (Preprint cond-mat/9301005)
Hsu T C and Dauriac J C A 1993 *Phys. Rev. B* **47** 14291
- [18] Mehta M L 1991 *Random Matrices* 2nd edn (San Diego, CA: Academic)
- [19] A mathematica notebook with the formulae and a simple example with a random choice of the assignable parameters is available at <http://physics.ucsc.edu/~sriram/demo-shastry.nb>
- [20] Pechukas P 1983 *Phys. Rev. Lett.* **51** 943
Yukawa T 1985 *Phys. Rev. Lett.* **54** 1883